

AD-A087 628

VIRGINIA UNIV CHARLOTTESVILLE SCHOOL OF MEDICINE F/G 6/5
THE EFFECTS OF ACUTE ALTERATIONS IN HEMODYNAMICS, OXYGEN AVAILA--ETC(U)
MAR 80 W P RITCHIE DAMD17-74-C-4014
NL

UNCLASSIFIED

1 of 1
Page 1 of 1




END

DATE

FILED

9-80

DTIC



ADA 087628

LEVEL

AD

THE EFFECTS OF ACUTE ALTERATIONS IN HEMODYNAMICS,
OXYGEN AVAILABILITY AND ACID-BASE BALANCE ON
THE PERMEABILITY OF THE GASTRIC MUCOSA

ANNUAL PROGRESS REPORT
(FOR THE PERIOD 1 OCT. 78 to 30 SEPT. 79)

DATE OF REPORT MARCH 1980

BY

WALLACE P. RITCHIE, JR., M.D.

SUPPORTED BY
U.S. ARMY MEDICAL RESEARCH AND DEVELOPMENT COMMAND
FORT DETRICK, FREDERICK, MARYLAND 21701

CONTRACT NO. DAMD 17-74-C-4014

UNIVERSITY OF VIRGINIA SCHOOL OF MEDICINE
CHARLOTTESVILLE, VIRGINIA 22908

DTIC
ELECTE
AUG 7 1980
C

APPROVED FOR PUBLIC RELEASE: DISTRIBUTION UNLIMITED

THE FINDINGS IN THIS REPORT ARE NOT TO BE CONSTRUED AS AN
OFFICIAL DEPARTMENT OF THE ARMY POSITION UNLESS SO DESIGNATED
BY OTHER AUTHORIZED DOCUMENTS.

DDC FILE COPY

80 7 31 066

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
	AD-A087628	9
4. TITLE (and Subtitle)	5. TYPE OF REPORT & PERIOD COVERED	
6 The Effects of Acute Alterations in Hemodynamics, Oxygen Availability, and Acid-Base Balance on the Permeability of the Gastric Mucosa.	Annual Progress Report. 1 Oct 1978-30 Sep 1979	
7. AUTHOR(s)	8. CONTRACT OR GRANT NUMBER(s)	
10 Wallace P. Ritchie, Jr., M.D., Ph.D.	15 DAMD 17-74-C-4014	
9. PERFORMING ORGANIZATION NAME AND ADDRESS	10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS	
University of Virginia School of Medicine Charlottesville, Virginia 22901	61102A 16 3A161102B71R 01.073 17 01	
11. CONTROLLING OFFICE NAME AND ADDRESS	12. REPORT DATE	
U.S. Army Medical Research & Development Command Frederick, Maryland 21701	11 March 1980	
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)	13. NUMBER OF PAGES	
12 8	8	
	15. SECURITY CLASS. (of this report)	
	Unclassified	
	15a. DECLASSIFICATION/DOWNGRADING SCHEDULE	
16. DISTRIBUTION STATEMENT (of this Report)		
Approved for public release, distribution unlimited		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number)		
Blood flow, ischemia, permeability, hydrochloric acid, histamine receptors, primary bile acids, secondary bile acids, ulcerogenesis.		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number)		
Using a previously described model for acute gastric mucosal ulcerogenesis, developed under the auspices of the current contract (Contract 68-699, 1975), studies from this laboratory during the period covered by this progress report indicate (1) that, in the face of an imposed topical acid load, H ₁ and H ₂ receptor blockade, either alone or in combination, effects no change in gastric mucosal permeability to cations or in mucosal blood flow and affords no protection against bile acid-ischemia induced ulcerogenesis; (2) that the addition of exogenous histamine neither protects nor augments lesion formation under		

#20 Continued:

these circumstances; (3) that the dihydroxy secondary bile acids are more damaging to gastric mucosa than are the trihydroxy primary bile acids; and (4) that in non-ischemic bile acid-treated gastric mucosa, mucosal blood flow increases in proportion to the magnitude of H⁺ back-diffusion induced, a response not mediated by histamine receptors.

Accession For	
NTIS GRA&I	<input checked="checked" type="checkbox"/>
DDC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution /	
Availability Codes	
Dist	Availability
A	

PROGRESS REPORT: CONTRACT DAMD 17-74-C-4014

I. TITLE OF RESEARCH CONTRACT:

The Effects of Acute Alterations in Hemodynamics, Oxygen Availability, and Acid-Base Balance on the Permeability of the Gastric Mucosa.

II. PRINCIPAL INVESTIGATOR:

Wallace P. Ritchie, Jr., M.D., Ph.D.
Department of Surgery
University of Virginia School of Medicine
Charlottesville, Virginia 22908
472-36-6392

III. PERIOD COVERED: 1 October 1978 to 30 September 1979

IV. PROGRESS REPORT:

(1) Effect of H_1 and H_2 receptor blockade on gastric mucosal permeability to cations and bile acid-induced ulcerogenesis. Recent reports suggest that, in the presence of topical acid, combined H_1 and H_2 receptor blockade decreases gastric mucosal permeability and H_2 blockade alone ameliorates bile-acid-induced "stress ulcer". The present study was undertaken to re-examine these assertions using the sensitive model outlined. Ex-vivo chambered wedges of mucosa were prepared in four groups of animals, each of which was studied during nine sequential 15 minute periods. Group A mucosa were exposed, during 1-3, to topical acid test solution alone (ATS), during 4-6, to ATS+2mM Na taurocholate (TC), and during 7-9 to ATS+TC+wedge-specific vasopressin (VP), 5×10^{-3} U/Kg-min. Groups B, C, and D were treated analogously except that group B was pretreated with cimetidine, 300mgm IV, group C with pyribenzamine, 10mgm/Kg IM, and group D, with both. The dose of cimetidine employed has been shown to produce significant inhibition of histamine stimulated gastric acid production in this model for a time period comparable to that of the current experiment. The dose of the H_1 receptor blocking agent was precisely the same as that reported to reduce, in combination with cimetidine, mucosal permeability to cations.

The parameters evaluated included (1) net flux of H^+ and Na^+ (μ Eq); (2) the potential difference (mV); (3) aminopyrine clearance (ml/min); and (4) the lesion index, graded 0-5. The results per 15 minutes (\pm SEM) indicate that, in mucosa exposed to ATS, $\Delta H^+ = -43 \pm 12$, $\Delta Na^+ = +133 \pm 13$, $PD = -63 \pm 1$, $AC = 2.3 \pm 0.4$, and $LI = 0$. During ATS+TC, $\Delta H^+ = -219 \pm 21$, $\Delta Na^+ = +206 \pm 17$, $PD = -45 \pm 1$, $AC = 2.8 \pm 0.5$, and $LI = 0$. H_1 , H_2 and $H_1 + H_2$ blockade effected no significant change under either circumstance. In bile acid treated ischemia mucosa;

GROUP	ΔH^+	ΔNa^+	\overline{PD}	AC	LI
Control (n=7)	-233 \pm 20	+323 \pm 20	-24 \pm 1	1.1 \pm 0.1	3.3 \pm 0.4
H ₁ (n=6)	-233 \pm 27	+333 \pm 24	-20 \pm 2	0.9 \pm 0.2	3.7 \pm 0.2
H ₂ (n=5)	-266 \pm 32	+361 \pm 22	-24 \pm 1	1.2 \pm 0.1	3.2 \pm 0.4
H ₁ +H ₂ (n=6)	-277 \pm 24	+324 \pm 17	-21 \pm 1	1.0 \pm 0.1	3.5 \pm 0.2

Thus, in the face of an imposed topical acid load, H₁ and H₂ blockade, both alone and in combination, effects no change in gastric mucosal permeability to cations or blood flow and affords no protection against bile acid-induced lesion formation.

(2) Effect of histamine stimulation on the magnitude of acute gastric mucosal damage induced by bile acids and ischemia.

It has been demonstrated in-vitro that burimamide-inhibited amphibian gastric mucosa is less resistant to bile acid induced damage, as judged by electrical indicies, than is spontaneously secreting mucosa. The present study used ex-vivo vascularized chambered wedges of proximal canine mucosa to examine the converse: the potential protective effect of active H⁺ secretion induced by histamine on acute mucosal ulcerogenesis caused by the topical application of Na taurocholate in acid solution and concomitant pharmacologic ischemia. Methods: Groups of dogs (n=5-6ea) were studied during 3 consecutive 30 minute periods. Group A= topical acid test solution (ATS) during periods (1), (2), (3). Group B= (1) ATS, (2) ATS, (3) ATS+vasopressin (VP), 0.1x10⁻²U/Kg-min delivered via the splenic artery. Group C= (1) ATS, (2) ATS+H⁺TC, (3) ATS+H⁺TC+VP. Parameters evaluated during each period=net H⁺ flux, aminopyrine clearance, and lesion index, graded 0-5. Results: In the absence of VP, TC significantly increased net H⁺ loss and AC relative to ATS; H resulted in significant H⁺ gain and further increased AC relative to ATS+TC; no lesions were noted under any circumstance. The results observed during period 3 (\pm SEM/30min):

	ATS	ATS+VP	ATS+TC+VP	ATS+H ⁺ TC+VP
ΔH^+ (μ Eq)	-109 \pm 43*	-108 \pm 60*	-242 \pm 53	-55 \pm 57*
AC (ml/min)	1.57 \pm 0.27	0.70 \pm 0.16*	1.13 \pm 0.27	1.57 \pm 0.49
LI (0-5)	0*	0.3 \pm 0.2*	2.4 \pm 0.3	1.6 \pm 0.5

*Significant difference vs. ATS+TC+VP.

Thus, the combination of topical bile acid in acid solution and relative mucosal ischemia (compared to ATS+TC, where AC=3.36 \pm 0.69ml/min) is acutely ulcerogenic. Concomitant histamine infusion neither protects nor augments lesion formation under these experimental conditions.

(3) Mediators of Bile Acid-Induced Enhancement of Gastric Mucosal Blood Flow. The topical application of bile acid in acid solution to non-ischemic gastric mucosa (GM) increases mucosal blood flow (MBF), an important cytoprotective phenomenon since, when it alone is blunted, mucosal ulcerogenesis results. The present study addressed the hypothesis that the magnitude of this response is related to the absolute amount of H⁺ "back diffusion" induced and that endogenous histamine release is the intramucosal mediator. Using vascularized chambered ex-vivo wedges of GM, 3 groups of dogs were studied during 3 sequential 30 minute periods. Group A (n=6): topical neutral test

solution (NTS; pH=7.0) during 1, NTS+2.5 mM sodium taurocholate (TC) during 2, and during 3, NTS+5 mM TC. Group B (n=11): topical acid test solution (ATS=100 mM HCl) during 1, ATS=2.5 TC during 2, and ATS=5 TC during 3. Group C (n=7) was treated in the same manner as group B except that each dog was pretreated with the H₁ and H₂ blockers, pyribenzamine (10 mg/Kg IM), and cimetidine (300 mg IV). Parameters evaluated: (1) net flux H⁺, Na⁺ (Δ H⁺, Δ Na⁺), (2) MBF, determine by radiolabelled microsphere embolization, (3) potential difference (PD). Results (\pm SEM)/30 minutes:

	NTS		ATS		NTS		ATS		NTS		ATS	
	0		-96 \pm 22*		0		-405 \pm 40**		0		-683 \pm 37**	
Δ H ⁺ (μ Eq)												
PD (mV)	-59 \pm 2		-62 \pm 1		-53 \pm 2*		-47 \pm 1**		-45 \pm 2*		-38 \pm 1**	
MBF (ml/min)	4.7 \pm 0.8		6.0 \pm 0.7		4.4 \pm 0.8		8.4 \pm 1.0**		4.2 \pm 1.0		10.4 \pm 1.0**	

*Significant difference vs NTS; ** Significant difference vs 0 nM TC, same group.

The addition of blockers effected no change from ATS alone in any parameter at any {TC} except at 5 TC, where MBF increased (to 15.8 \pm 2.6 ml/min.). CONCLUSIONS: (1) In non-ischemic GM, MBF increases in the presence of greater than normal Δ H⁺. (2) The amplitude of this response is dependent on the absolute magnitude of the increase in H⁺ "back diffusion" observed (p<0.01). (3) Enhanced MBF is not mediated by H₁ and H₂ receptor related histamine release. In fact, receptor blockage augments the response at 5mM TC.

(4) Ulcerogenic Potential of Primary vs. Secondary Bile Acids in Gastric Mucosa. Although the secondary (2^o) bile acids are more damaging to colonic mucosa than are the primary (1^o) bile acids, the existence of a similar circumstance in gastric mucosa has been inadequately documented. The present study addressed this problem by comparing the effect of an identical (and physiologic) concentration of the 2^o BA, taurodeoxycholic a. (TDC) to its parent 1^o BA taurocholic a. (TC) in both ischemic and non-ischemic GM. METHODS: Using chamber ex-vivo wedges of proximal GM, 2 groups of dogs were studied during 9 sequential 15 minute periods. Group A (4 dogs)=topical acid test solution alone (ATS=100mM HCl) during periods 1-3; ATS+1mM TC during 4-6; and, during 7-9, ATS+1mM TC+vasopressin (VP=5x10⁻³U/Kg-min) infused into the splenic artery. Group B (4 dogs) were similarly treated except that the BA employed was TDC. Parameters evaluated=(1) net flux H⁺, Na⁺ (Δ H⁺, Δ Na⁺), (2) potential difference (PD), (3) mucosal blood flow (MBF), determined by radiolabeled microsphere embolization, and (4) the lesion index (LI), graded 0-5 by an independent observer using photographs. RESULTS (\pm SEM)/15 minutes:

GROUP	ATS (1-3)		ATS+1mM BA (4-6)		ATS+1mM BA+VP (7-9)	
	A	B	A	B	A	B
PD (mV)	-64 \pm 1	-63 \pm 1	-60 \pm 1*	-48 \pm 1**	-41 \pm 1*	-28 \pm 1**
Δ H ⁺ (μ Eq)	-8 \pm 15	-1 \pm 18	-97 \pm 13*	-157 \pm 20**	-105 \pm 21*	-250 \pm 30**
MBF (ml/min)	11.2 \pm 2.0	12.0 \pm 2.5	9.0 \pm 1.4	13.6 \pm 4.8	2.5 \pm 0.7*	2.9 \pm 0.5*
LI (0-5)	0	0	0	0	0.9 \pm 0.4*	2.6 \pm 0.4*

+P 0.05 vs. ATS, same group; *p 0.025 vs. A(TC)

Thus, at an identical (and physiologic) concentration, 2° BA induce a greater depression on PD, a more profound increase in mucosal permeability to H^{+} , and in ischemic GM, a more severe degree of gross mucosal damage than do 1° BA. CONCLUSION: 2° BA are more damaging to GM than are 1° BA, an effect undoubtedly related to their greater lipid solubility and consequent capacity to disrupt cell membranes.

V. PUBLICATIONS DURING THE CONTRACT YEAR:

1. Ritchie, W.P., Jr. and Cherry, K.J., Jr.: Influence of $[H^{+}]$ on Bile Acid Induced Acute Gastric Mucosal Ulcerogenesis. Ann. Surg. 189:637-642, 1979.
2. Ritchie, W.P., Jr. and McRae, D.B., Jr.: Effect of H_1 and H_2 Receptor Blockade on Gastric Mucosal Permeability to Cations and Bile Acid Induced Ulcerogenesis. Gastroent. 76:1227, 1979. Abstract.
3. Ritchie, W.P., Jr., McRae, D.B., Jr. and Felger, T.S.: A-V Acid-Base Balance During Acute Gastric Mucosal Ulcerogenesis. Am. J. Surg. 139:22-26, 1980.
4. Ritchie, W.P., Jr.: Invited Commentary on "Restraint Induced Stress Ulcer. Parts I and II" World Journal of Surgery 4:101-103, 1980.
5. Felger, T.S., and Ritchie, W.P., Jr.: Ulcerogenic Potential of 1° vs. 2° Bile Acids in Gastric Mucosa. Surgical Forum 30:324-326, 1979.

DISTRIBUTION LIST

4 copies

HQDA (SGRD-SI)
Fort Detrick
Frederick, MD. 21701

12 copies

Defense Technical Information Center (DTIC)
ATTN: DTIC-DDA
Cameron Station
Alexandria, Virginia 22314

1 copy

Dean
School of Medicine
Uniformed Services University of the
Health Sciences
4301 Jones Bridge Road
Bethesda, Maryland 20014

1 copy

Superintendent
Academy of Health Sciences, US Army
ATTN: AHS-COM
Fort Sam Houston, Texas 78234

4 copies

Commander
Letterman Army Institute
of Research (LAIR) Bldg. 1110
ATTN: Dr. J. Ryan Neville
Presidio of San Francisco, CA 94129